



Collagen, Human Type VI

<u>Cat. No.</u>	<u>Form</u>	<u>Quantity</u>
1290-01S	Purified protein	0.25 mg

DESCRIPTION

Source Placental villi
Purification Controlled and limited pepsin digestion, followed by selective salt precipitation.
Purity >90% as judged by SDS-PAGE analysis

RESEARCH APPLICATIONS

- Type VI collagen standard
- Antigen for antibody production
- Coating material for cell culture studies
- Formation of collagen gels

CHARACTERIZATION

To insure lot-to-lot consistency, each batch of product is tested to conform to characteristics of a standard reference reagent using SDS-PAGE analysis

HANDLING AND STORAGE

- The purified protein is supplied as a solution of 0.25 mg collagen in 0.5 mL of 500 mM acetic acid (C = 0.5 mg/mL).
- The collagen may be transferred into physiological buffers by dialysis at 2-8°C.
- Purified collagen may be stored at 2-8°C for several months without appreciable loss of activity. For long-term storage, the purified collagen may be aliquoted and stored at -20°C. **Avoid repeated freezing and thawing.**

REFERENCES

1. Miller, E.J. and R.K. Rhodes. 1982. *Meth. Enzymol.* 82:33.
2. Miller, E.J. In: *Extracellular Matrix Biochemistry*, K.A. Piez and A.H. Reddi, eds. Elsevier, New York, 1984. pp. 41-81.

For Research Use Only. Not for Diagnostic or Therapeutic Use.